

A class of integration by parts formulae in stochastic analysis I

K. D. Elworthy and Xue-Mei Li
Mathematics Institute
University of Warwick
Coventry CV4 7AL, U.K.

1 Introduction

Consider a Stratonovich stochastic differential equation

$$dx_t = X(x_t) \circ dB_t + A(x_t)dt \quad (1)$$

with C^∞ coefficients on a compact Riemannian manifold M , with associated differential generator $\mathcal{A} = \frac{1}{2}\Delta_M + Z$ and solution flow $\{\xi_t : t \geq 0\}$ of random smooth diffeomorphisms of M . Let $T\xi_t : TM \rightarrow TM$ be the induced map on the tangent bundle of M obtained by differentiating ξ_t with respect to the initial point. Using an observation by A. Thalmaier we will extend the basic formula of [EL94] to obtain

$$\mathbb{E}dF(T\xi.(h.)) = \mathbb{E}F(\xi.(x)) \int_0^T \left\langle T\xi_s(\dot{h}_s), X(\xi_s(x)) dB_s \right\rangle \quad (2)$$

where $F \in \mathcal{F}C_b^\infty(C_x(M))$, the space of smooth cylindrical functions on the space $C_x(M)$ of continuous paths $\gamma : [0, T] \rightarrow M$ with $\gamma(0) = x$, dF is its derivative, and $h.$ is a suitable adapted process with sample paths in the Cameron-Martin space $L_0^{2,1}([0, T]; T_x M)$. Set $\mathcal{F}_t^x = \sigma\{\xi_s(x) : 0 \leq s \leq t\}$. Taking conditional expectation with respect to \mathcal{F}_T^x , formula (2) yields integration by parts formulae on $C_x(M)$ of the form

$$\mathbb{E}dF(\gamma)(\bar{V}^h) = \mathbb{E}F(\gamma)\delta\bar{V}^h(\gamma) \quad (3)$$

where \bar{V}^h is the vector field on $C_x(M)$

$$\bar{V}^h(\gamma)_t = \mathbb{E}\{T\xi_t(h_t) | \xi.(x) = \gamma\}$$

and $\delta\bar{V}^h : C_x(M) \rightarrow \mathbb{R}$ is given by

$$\delta\bar{V}^h(\gamma) = \mathbb{E} \left\{ \int_0^T \langle T\xi_s(\dot{h}_s), X(\xi_s(x))dB_s \rangle \mid \xi(\cdot) = \gamma \right\}.$$

When h is adapted to \mathcal{F}^x results from [ELJL95] extending [EY93] give explicit expressions for \bar{V}^h and $\delta\bar{V}^h$ in terms of the Ricci curvature of the LeJan-Watanabe connection associated to (1). Equation (3) then reduces to a Driver's integration by parts formula, Theorem 3.3 below, but no hypothesis of torsion skew symmetry of the connection is required: the integration by parts formulae follow for the adjoint of any metric connection. In particular for any such connection there is a Hilbert "tangent space" of "good" directions obtained by parallel translation of the Cameron-Martin space of paths in T_xM . (In fact it is the "Ricci flow" or "Dohrn-Guerra parallel translation" (see Nelson [Nel84]), leading to the "damped gradient" ([FM93]) which occurs more naturally). However, in Remark 2.4, we show that in this case \bar{V}_h is in the class for which integration by parts formulae are known, so that the results of 2.3, 3.3, 3.5 are not claimed to be new in substance.

Although this filtering out of the extraneous noise gives intrinsic results comparable to those of Driver [Dri92], this viewpoint throws away a lot of the structure we have. Moreover integration by parts formulae such as (2) should have some connection with quasi-invariance properties of flows associated to the vector fields. Flows for the \bar{V}^h on $C_x(M)$ do not appear to be easy to analyse in general. However in §3 we show that in the context of Diff M valued processes there are very natural flows associated and (2) has a rather natural geometric interpretation. This leads to another elementary proof of (2) and in Theorem 4.1 we use this method to obtain integration by parts formulae for the free path space.

There are at least 3 proofs of (2). The first given here is via Itô's formula and elementary martingale calculus (it requires F to be cylindrical), the second given here is based on the Girsanov-Maruyama theorem (and works for more general F), and a third method would be to deduce it from the standard integration by parts formula on Wiener space applied to the functional $F \circ \xi$. Indeed this work was stimulated by D. Bell and D. Nualart pointing out that this third approach could be used to deduce the basic formula of [EL94]. The point made (and carried out) in [Elw92] and [EL94] that the first approach can be applied directly to 'Ricci flows' instead of derivative flows to give intrinsic formulae without stochastic flows, also needs to be emphasized: see also [SZ].

There are also now many proofs of Driver's results for $C_x(M)$ and for the free path space and their extensions. See [Hsu95], [ES95], [LN] (with a very concise proof), [AM], [Aid], and [CM].

Acknowledgment: This research was supported by SERC grant GR/H67263 and stimulated and helped by our contacts with A. Thalmaier.

2 The integration by parts formula from finite dimensional manifolds to path spaces

In this section we deduce by induction an integration by parts formula on the path space from a formula on the base manifold M . The key is to obtain formula (10) for M .

Let $h : \Omega \times [0, T] \rightarrow T_x M$ be an adapted process with $h(\omega) : [0, T] \rightarrow T_x M$ in $L^{2,1}$ for almost all ω .

Lemma 2.1 *If $h : \Omega \times [0, T] \rightarrow T_x M$ is adapted, $L^{2,1}$ for a.s. ω and $\left(\int_0^T |\dot{h}_s|^2 ds\right)^{1/2} \in L^{1+\epsilon}$ for some $\epsilon > 0$. Then for $t < T$,*

$$\begin{aligned} & \mathbb{E} \left\{ \int_0^t \langle T\xi_s(\dot{h}_s), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\} \\ &= \mathbb{E} \left\{ \int_t^T \langle T\xi_s(-), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\}. \end{aligned} \quad (4)$$

If furthermore h is non-random then for $t \leq T$,

$$\begin{aligned} & \mathbb{E} \left\{ \int_0^t \langle T\xi_s(\dot{h}_s), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\} \\ &= \mathbb{E} \left\{ \int_0^t \langle T\xi_s(-), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\}. \end{aligned} \quad (5)$$

Proof. First by the Burkholder-Davis-Gundy inequality, for some constant c_1 ,

$$\begin{aligned} & \mathbb{E} \left| \int_0^T \langle T\xi_s(\dot{h}_s), X(\xi_s(x)) dB_s \rangle \right| \leq c_1 \mathbb{E} \left(\int_0^T |T\xi_s(\dot{h}_s)|^2 ds \right)^{\frac{1}{2}} \\ & \leq c_1 \left(\mathbb{E} \sup_{0 \leq s \leq T} |T_x \xi_s|^{\frac{1+\epsilon}{\epsilon}} \right)^{\frac{\epsilon}{1+\epsilon}} \left[\mathbb{E} \left(\int_0^T |\dot{h}_s|^2 ds \right)^{\frac{1+\epsilon}{2}} \right]^{\frac{1}{1+\epsilon}}. \end{aligned}$$

This is finite since $\sup_{0 \leq s \leq t} |T_x \xi_s| \in L^q$ for all $1 \leq q < \infty$, e.g. see [Li94]. Moreover, since the adapted processes in $L^\infty(\Omega, \mathcal{F}, \mathbb{P}; C^1([0, T]; T_x M))$ are

dense in the subspace of adapted processes in $L^{1+\epsilon}(\Omega, \mathcal{F}, \mathbb{P}; L^{2,1}([0, T]; T_x M))$, this estimate allows us to assume that h belongs to the former space.

Set $M_t = \int_0^t \langle T_x \xi_s(-), X(\xi_s(x)) dB_s \rangle$. Then $\{M_t\}$ is a $T_x^* M$ valued local martingale. If $0 = t_0 < t_1 < \dots < t_l = t$ is a partition of $[0, t]$, $\Delta_j t = t_{j+1} - t_j$, and $\Delta_j M = M_{t_{j+1}} - M_{t_j}$, then

$$\sum_{j=1}^{l-1} \Delta_j M(\dot{h}_{t_j}) \rightarrow \int_0^t \dot{h}_s dM_s = \int_0^t \langle T \xi_s(\dot{h}_s), X(\xi_s(x)) dB_s \rangle \quad (6)$$

and the convergence is in L^1 .

On the other hand if $v_0 \in T_x M$ and P_t is the probabilistic semigroup associated to the S.D.E. and f a bounded measurable function then

$$d(P_t f)(v_0) = \frac{1}{T} \mathbb{E} f(\xi_T(x)) \int_0^T \langle T \xi_s(v_0), X(\xi_s(x)) dB_s \rangle. \quad (7)$$

See [EL94]. However by an observation of Thalmaier: the same proof shows that for any $r, h \in [0, T]$ with $h > 0$ and $r + h \leq T$

$$d(P_t f)(v_0) = \frac{1}{h} \mathbb{E} f(\xi_T(x)) \int_r^{r+h} \langle T \xi_s(v_0), X(\xi_s(x)) dB_s \rangle$$

c.f. [SZ]. From these two formulae we obtain:

$$\begin{aligned} & \mathbb{E} \left\{ \frac{1}{T} \int_0^T \langle T \xi_s(v_0), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\} \\ &= \mathbb{E} \left\{ \frac{1}{h} \int_r^{r+h} \langle T \xi_s(v_0), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\}. \end{aligned} \quad (8)$$

For any $0 \leq r \leq T$, let $\{\xi_s^r(x) : r \leq s \leq T, x \in M\}$ be the solution flow to (1) starting from x at time r . The flow ξ^r can be taken to be adapted to a filtration $\{\mathcal{F}_s^r : r \leq s \leq T\}$ independent of \mathcal{F}_r , and then we have $\xi_s^r \xi_r = \xi_s$, almost surely, $r \leq s \leq T$. From this, time homogeneity, and (8),

$$\begin{aligned} & \mathbb{E} \left\{ \sum_{j=1}^{l-1} \Delta_j M(\dot{h}_{t_j}) \mid \xi_T(x) \right\} \\ &= \mathbb{E} \left\{ \sum_{j=1}^{l-1} \Delta_j t \frac{1}{\Delta_j t} \int_{t_j}^{t_{j+1}} \langle T \xi_s^{t_j} (T \xi_{t_j}(\dot{h}_{t_j})), X(\xi_s^{t_j}(\xi_{t_j}(x))) dB_s \rangle \mid \xi_T^{t_j}(\xi_{t_j}(x)) \right\} \\ &= \mathbb{E} \left\{ \sum_{j=1}^{l-1} \Delta_j t \frac{1}{T-t} \int_t^T \langle T \xi_s^{t_j} (T \xi_{t_j}(\dot{h}_{t_j})), X(\xi_s^{t_j}(\xi_{t_j}(x))) dB_s \rangle \mid \xi_T^{t_j}(\xi_{t_j}(x)) \right\} \end{aligned}$$

$$\begin{aligned}
&= \mathbb{E} \left\{ \sum_{j=1}^{l-1} \Delta_j t \frac{1}{T-t} \int_t^T \langle T\xi_s(\dot{h}_{t_j}), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\} \\
&\rightarrow \mathbb{E} \left\{ \int_t^T \langle T\xi_s(-), X(\xi_s(x)) dB_s \rangle > \frac{h_t - h_0}{T-t} \mid \xi_T(x) \right\}.
\end{aligned}$$

Comparing with (6) this gives the first required identity. When h is non-random the second follows immediately from (8). \blacksquare

Remark:

As in [SZ] a further modification is possible replacing (8) by:

$$\begin{aligned}
&\frac{1}{T} \mathbb{E} \left\{ \int_0^T \langle T\xi_s(v_0), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\} \\
&= \frac{1}{\int_0^T \Psi(r) dr} \mathbb{E} \left\{ \int_0^T \Psi(s) \langle T\xi_s(v_0), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\}
\end{aligned}$$

for $\Psi : [0, T] \rightarrow \mathbb{R}$ integrable with $\int_0^T \Psi(r) dr \neq 0$. The argument leads to, for non-random h ,

$$\begin{aligned}
&\mathbb{E} \left\{ \int_0^t \langle T\xi_s(\dot{h}_s), X(\xi_s(x)) dB_s \rangle \mid \xi_T(x) \right\} \\
&= \mathbb{E} \left\{ \int_0^T \Psi(s) \langle T\xi_s(-), X(\xi_s(x)) dB_s \rangle > \left(\frac{h_t - h_0}{\int_0^T \Psi(r) dr} \right) \mid \xi_T(x) \right\}. \tag{9}
\end{aligned}$$

Corollary 2.2 *Under the conditions of the lemma, for any C^1 function $f : M \rightarrow \mathbb{R}$,*

$$\mathbb{E} f(\xi_T(x)) \int_0^T \langle T\xi_s(\dot{h}_s), X(\xi_s(x)) dB_s \rangle = \mathbb{E} df(T\xi_T(h_T - h_0)). \tag{10}$$

Proof. First by the composition property of solution flows,

$$\begin{aligned}
&\mathbb{E} \left\{ \int_t^T \langle T\xi_s(-), X(\xi_s(x)) dB_s \rangle > \frac{h_t - h_0}{T-t} \mid \xi_T(x) \right\} \\
&= \mathbb{E} \left\{ \int_t^T \langle T\xi_s^t(-), X(\xi_s^t(\xi_t(x))) dB_s \rangle > \frac{T\xi_t(h_t - h_0)}{T-t} \mid \xi_T^t(\xi_t(x)) \right\}.
\end{aligned}$$

As in the proof of the lemma, (4) yields

$$\begin{aligned}
& \mathbb{E}f(\xi_T(x)) \int_0^t \langle T\xi_s(\dot{h}_s), X(\xi_s(x))dB_s \rangle \\
&= \mathbb{E}f(\xi_T^t(\xi_t(x))) \int_t^T \langle T\xi_s^t(-), X(\xi_s^t(\xi_t(x)))dB_s \rangle \frac{T\xi_t(h_t - h_0)}{T - t} \\
&= \mathbb{E}\{dP_{T-t}(f)(T\xi_t(h_t - h_0))\}
\end{aligned}$$

by [EL94], since \mathcal{F}^t is independent of \mathcal{F}_t . Now let t increase to T and the required result follows. \blacksquare

Next consider a cylindrical function F on $C_x(M)$, the space of continuous paths with base point x . Write

$$F(\gamma.) = f(\gamma_{t_1}, \dots, \gamma_{t_k}),$$

for $(t_1, \dots, t_k) \in [0, T]^k$, $\gamma \in C_x(M)$ and f a smooth function on M^k . Suppose $h_0 = 0$ and consider the tangent vector field $V^h(\xi.(x))$ along $\{\xi_t(x) : 0 \leq t \leq T\}$ on $C_x(M)$ given by

$$V^h(\xi.)_t = T_x\xi_t(h_t).$$

Then

$$dF(V^h(\xi.)) = \sum_{j=1}^k d^j f_{\xi_{t_j}}(V^h(\xi.)_{t_j}). \quad (11)$$

Here $\xi_{t_j} = (\xi_{t_1}, \dots, \xi_{t_k})$ and $d^j f$ is the partial derivative of f in the j th direction.

Let

$$\delta V^h(\xi.) = \int_0^T \langle T_x\xi_s(\dot{h}_s), X(\xi_s(x))dB_s \rangle.$$

Theorem 2.3 *Let $h : [0, T] \times \Omega \rightarrow T_x M$ be an adapted stochastic process with almost surely all $h(\omega) \in L_0^{2,1}$ and $\mathbb{E}\left(\int_0^T |\dot{h}_s|^2 ds\right)^{\frac{1+\epsilon}{2}} < \infty$ for some $\epsilon > 0$.*

Then

$$\mathbb{E}dF(V^h(\xi.)) = \mathbb{E}F(\xi.(x))\delta V^h(\xi.). \quad (12)$$

Proof. We prove by induction on k . When $k = 1$, this is just (10), the formula for functions. Let $\Omega = C_0([0, T]; \mathbb{R}^n)$ be the canonical probability space. We set $\Omega_1 = C_0([0, t_1]; \mathbb{R}^n)$ and $\Omega_2 = C_0([t_1, T]; \mathbb{R}^n)$. There is then the standard decomposition of filtered spaces

$$\begin{aligned}
& \{\Omega, \mathcal{F}, \mathcal{F}_t, 0 \leq t \leq T, \mathbb{P}\} \\
&= \{\Omega_1, \mathcal{F}, \mathcal{F}_t, 0 \leq t \leq t_1, \mathbb{P}_{\neq}\} \times \{\Omega_2, \mathcal{F}, \mathcal{F}_t^{t_1}, t_1 \leq t \leq T, \mathbb{P}_{\neq}\}
\end{aligned}$$

in the sense that $\mathcal{F}_t = \mathcal{F}_t * \Omega_2$ if $t \leq t_1$, and $\mathcal{F}_t = \mathcal{F}_{t_1} * \mathcal{F}_t^{t_1}$ if $t \geq t_1$. As before let $\xi_t^{t_1}(y_0), t_1 \leq t \leq T, y_0 \in M$ be the solution flow to (1) starting at time t_1 , i.e. $\xi_{t_1}^{t_1}(y_0) = y_0$. We will consider it as a function of $\omega_2 \in \Omega_2$, adapted to \mathcal{F}^{t_1} , while $\{\xi_t : 0 \leq t \leq t_1\}$ will be considered on Ω_1 , and $\{\xi_t : t_1 \leq t \leq T\}$ on $\Omega_1 \times \Omega_2 = \Omega$. The composition property for flows gives

$$\xi_t^{t_1}(\xi_{t_1}(x, \omega_1), \omega_2) = \xi_t(x, (\omega_1, \omega_2)), \quad \text{each } t_1 \leq t \leq T, a.s.$$

Assume the required result holds for cylindrical functions depending on $k - 1$ times, some $k \in \{2, 3, \dots\}$. Take $y_0 \in M$ and define $f_1^{y_0} : M^{k-1} \rightarrow \mathbb{R}$ and $F_1^{y_0} : \Omega_2 \rightarrow \mathbb{R}$ by:

$$f_1^{y_0}(x_1, \dots, x_{k-1}) = f(y_0, x_1, \dots, x_{k-1})$$

and

$$F_1^{y_0}(\omega_2) = f(y_0, \xi_{t_2}^{t_1}(y_0, \omega_2), \dots, \xi_{t_k}^{t_1}(y_0, \omega_2)).$$

Take $h^1 : \Omega_2 \rightarrow L_0^{2,1}([t_1, T]; T_{y_0}M)$, adapted to \mathcal{F}^{t_1} , and with $\mathbb{E} \left(\int_{t_1}^T |h_s^1|^2 ds \right)^{\frac{1+\epsilon}{2}}$ finite. By time homogeneity our inductive hypothesis gives

$$\begin{aligned} & \sum_{j=2}^k \int_{\Omega_2} d^j f(y_0, \xi_{t_2}^{t_1}(y_0, \omega_2), \dots, \xi_{t_k}^{t_1}(y_0, \omega_2)) \left(T \xi_{t_j}^{t_1}(h_{t_j}^1(\omega_2), \omega_2) \right) d\mathbb{P}_{\neq}(\omega_2) \\ &= \int_{\Omega_2} f(y_0, \xi_{t_2}^{t_1}(y_0, \omega_2), \dots, \xi_{t_k}^{t_1}(y_0, \omega_2)) \times \\ & \quad \int_{t_1}^T \left\langle T \xi_r^{t_1}(\dot{h}_r^1(\omega_2), \omega_2), X(\xi_r^{t_1}(y_0, \omega_2)) dB_r(\omega_2) \right\rangle d\mathbb{P}_{\neq}(\omega_2). \end{aligned} \tag{13}$$

Now for $\omega_1 \in \Omega_1$ (outside of a certain measure zero set) we can take $y_0 = \xi_{t_1}(x_0, \omega_1)$ and

$$h_t^1(\omega_2) = T \xi_{t_1}(h_t(\omega_1, \omega_2) - h_{t_1}(\omega_1), \omega_2).$$

Then, for almost all $\omega_1 \in \Omega_1$, we have h^1 adapted to \mathcal{F}^{t_1} . Substitute this in (13). Using the composition property, and then integrating over Ω_1 yields

$$\begin{aligned} & \sum_{j=2}^k \mathbb{E} d^j f(\xi_{t_j}) \left(T \xi_{t_j}(h_{t_j} - h_{t_1}) \right) \\ &= \mathbb{E} f(\xi_{t_1}(x)) \int_{t_1}^T \left\langle T \xi_r(\dot{h}_r), X(\xi_r(x)) dB_r \right\rangle. \end{aligned} \tag{14}$$

On the other hand we can define $g : M \rightarrow \mathbb{R}^1$ by

$$g(x) = \int_{\Omega_2} f(x, \xi_{t_2}^{t_1}(x, \omega_2), \dots, \xi_{t_k}^{t_1}(x, \omega_2))$$

and apply formula (10) to g to obtain:

$$\int_{\Omega_1} dg(T\xi_{t_1}(h_{t_1}))d\mathbb{P}_{\mu^*}(\omega_1) = \int_{\Omega_1} g(\xi_{t_1}(x)) \int_0^{t_1} \langle T\xi_r(\dot{h}_r), X(\xi_r(x))dB_r \rangle d\mathbb{P}_{\mu^*}(\omega_1).$$

But note that

$$\int_{\Omega_1} dg(T\xi_{t_1}(h_{t_1}))d\mathbb{P}_{\mu^*}(\omega_1) = \sum_{j=1}^k \mathbb{E}d^j f_{\xi_{\underline{t}}}(T\xi_{t_j}(h_{t_1}))d\mathbb{P}_{\mu^*}(\omega_1),$$

and therefore

$$\sum_{j=1}^k \mathbb{E}d^j f_{\xi_{\underline{t}}}(T\xi_{t_j}(h_{t_1})) = \mathbb{E}f(\xi_{\underline{t}}) \int_0^{t_1} \langle T\xi_r(\dot{h}_r), X(\xi_r(x))dB_r, \rangle \quad (15)$$

Adding (14) we arrive at (12):

$$\sum_{j=1}^k \mathbb{E}d^j f_{\xi_{\underline{t}}}(T\xi_{t_j}(h_{t_j})) = \mathbb{E}f(\xi_{\underline{t}}(x)) \int_0^T \langle T\xi_r(\dot{h}_r), X(\xi_r(x))dB_r \rangle.$$

■

B. Let $\tilde{\nabla}$ be a metric connection for the manifold M with torsion T , and $\tilde{\nabla}'$ its adjoint connection defined by

$$\tilde{\nabla}'_{V_1} V_2 = \tilde{\nabla}_{V_1} V_2 - T(V_1, V_2).$$

Here V_1, V_2 are vector fields. Let \tilde{R} be the curvature tensor of $\tilde{\nabla}$ and define $\tilde{Ric}^\# : TM \rightarrow TM$ by $\tilde{Ric}^\#(v) = \text{trace } \tilde{R}(v, -)$. If $\{x_s\}$ is a diffusion on M with generator $\frac{1}{2}\text{trace}\tilde{\nabla}\text{grad} + L_Z$ denote by $\tilde{\parallel}_s$ the parallel transport along $\{x_s\}$, and $\{\tilde{B}_s : 0 \leq s \leq t\}$ the martingale part of the anti-development of $\{x_s : 0 \leq s \leq t\}$ using $\tilde{\parallel}_s$, a Brownian motion on $T_{x_0}M$. Let $v_s = \tilde{W}_s^Z(v_0)$ be the solution to

$$\frac{\tilde{D}'}{\partial s} v_s = -\frac{1}{2}\tilde{Ric}^\#(v_s) + \tilde{\nabla}Z(v_s)$$

starting from $v_0 \in T_{x_0}M$. Here \tilde{D}' denotes the covariant differentiation along the paths of $\{x_t\}$ using the adjoint connection. We will show that (12) implies Driver's integration by parts formula. However we do not need to assume $\tilde{\nabla}'$ (or equivalently $\tilde{\nabla}$) is torsion skew symmetric.

Corollary 2.4 *Let F be a cylindrical function on $C_{x_0}(M)$. Suppose $h : [0, T] \times \Omega \rightarrow T_{x_0}M$ is adapted to the filtration of $\{x_s : 0 \leq s < \infty\}$ and such*

that $h(\omega)$ is in $L_0^{2,1}$ for almost all ω and $h \in L^{1+\epsilon}(\Omega, \mathcal{F}, \mathbb{P}; L_0^{2,1}([0, T]; T_{x_0}M))$ for some $\epsilon > 0$. Then

$$\mathbb{E}dF(\tilde{W}^Z(h.)) = \mathbb{E}F(\xi.(x_0)) \int_0^T \langle \tilde{W}_s^Z(\dot{h}_s), \tilde{\int}_s d\tilde{B}_s \rangle. \quad (16)$$

When $\tilde{\nabla}'$ is metric for some Riemannian metric on M , it suffices to have $h \in L^1(\Omega, \mathcal{F}, \mathbb{P}; L_0^{2,1}([0, T]))$.

Proof. By a result of [ELJL95] we can choose X such that $\tilde{\nabla}$ equals the Le Jan-Watanabe connection induced from the stochastic differential equation

$$dx_t = X(x_t) \circ dB_t + Z(x_t)dt$$

and the solution flow $\{\xi.(x)\}$ has generator $\frac{1}{2}\text{trace}\tilde{\nabla}\text{grad} + L_Z$ (c.f. Corollary 3.4 of [ELJL95]). Moreover the conditioned process of the derivative flow $T\xi_t(v_0)$ with respect to the natural filtration of $\{\xi.(x_0)\}$ is given by $\{\tilde{W}^Z(v_0)\}$:

$$\mathbb{E}\{T\xi_t(v_0) \mid \mathcal{F}_T^{x_0}\} = \tilde{W}_t^Z(v_0),$$

by Theorem 3.2 of [ELJL95] extending [EY93]. The result follows since \tilde{B}_t equals $\int_0^t \tilde{\int}_s^{-1} X(\xi_s(x_0))dB_s$.

If $\tilde{\nabla}'$ is metric for some Riemannian metric then $\sup_{0 \leq s \leq t} |\tilde{W}_s^Z|$ is in $L^\infty(\Omega, \mathcal{F}, \mathbb{P})$ and so the Burkholder-Davis-Gundy inequality used as in the proof of Lemma 2.1 allows us to take $\epsilon = 0$. \blacksquare

Remarks 2.5. (i). Let $S : TM \times TM \rightarrow TM$ be a tensor fields of type (1,2), and let ∇ refer to the Levi-Civita connection of M . Then, by [KN69] p.146, a connection $\tilde{\nabla}$ can be defined by

$$\tilde{\nabla}_{V_1}(V_2) = \nabla_{V_1}(V_2) + S(V_1, V_2)$$

for vector fields V_1, V_2 . and all linear connections on M can be obtained this way. It is easy to see that $\tilde{\nabla}$ is metric if and only if

$$\langle S(W, U), V \rangle = - \langle U, S(W, V) \rangle$$

for all vector fields U, V, W , i.e. if and only if $S(W, -)$ is skew symmetric. On the other hand the adjoint connection is given by

$$\tilde{\nabla}'_{V_1}(V_2) = \nabla_{V_1}(V_2) + S(V_2, V_1)$$

so that it is torsion skew symmetric if also $S(-, W)$ is skew symmetric. In terms of the Levi-Civita connection our vector fields \tilde{V}^h for which the integration by parts formula hold therefore satisfy an equation of the form

$$D\bar{V}_t^h = -S(\bar{V}_t^h, \circ dx_t) + \Lambda_t(\bar{V}_t^h)dt + W_t^h(\dot{h}_t)dt + \nabla A(\bar{v}_t^h)dt$$

where Λ_t is linear (also depending on S). In particular they are “tangent processes” in the sense proposed by Driver, for which integration by parts formulae are known: see [Dri95b], [CM], [AM], and [Aid], [Dri95a].

(ii) For cylinder functions depending on one time only such integration by parts formulae go back to Bismut [Bis84].

3 Geometric interpretation and a shorter proof

A. The processes $T_x\xi_t(h_t)$ cannot strictly speaking be considered as tangent vectors or vector fields on $C_x(M)$. In some sense they form tangent vectors at $\xi.(x, -)$ to the space of processes (or semi-martingales)

$$[0, T] \times \Omega \rightarrow M$$

since $T_x\xi_t(h_t(\omega), \omega) \in T_{\xi_t(x, \omega)}M$ for $(t, \omega) \in [0, T] \times \Omega$ or equivalently as ‘tangent vectors’ to the space of random variables

$$\Omega \rightarrow C_x(M)$$

at $\omega \mapsto \xi.(x, \omega)$. However c.f. [Dri92] there is still no natural associated flow. In fact the most natural interpretation takes into account the variable x and replaces $C_x(M)$ by $P_{id}\text{Diff}M$ the space of paths on the diffeomorphism group of M , as we now describe.

Let $\text{Diff}M$ be the space of C^∞ diffeomorphisms of M . We can consider it with a rather formal differential structure or if the reader prefers it can be replaced by a suitable Sobolev space of diffeomorphisms, to give a Hilbert manifold (as in [Elw82] following [EM70]). In any case the tangent space $T_\alpha(\text{Diff}M)$ will be identified with all vector fields on M over α i.e. smooth $v : M \rightarrow TM$ such that $v(x) \in T_{\alpha(x)}M$ for all $x \in M$. If $P\text{Diff}M$ refers to continuous paths $\phi : [0, T] \rightarrow \text{Diff}M$ with $\phi(0) = id_M$ then $T_\phi P\text{Diff}M$ will be identified with continuous $v : [0, T] \rightarrow T\text{Diff}M$ vanishing at $t = 0$, such that $v(t) \in T_{\phi(t)}\text{Diff}M$, or equivalently $v : [0, T] \times M \rightarrow TM$ with $v(t)(x) \in T_{\phi(t)(x)}M$.

B. Given our S.D.E. (1) now take $h \in L_0^{2,1}([0, T]; \mathbb{R}^n)$. There is X^h , the time dependent vector field $X(\cdot)(h_t)$ on M . From this we obtain a field U^h on $P\text{Diff}M$ by

$$U^h(\phi)_t(x) = T_x\phi_t(X(x)h_t). \tag{17}$$

This is just the left invariant vector field on $P\text{Diff}M$ corresponding to $X^h \in T_e P\text{Diff}M$ for $e(t) = id_M$, $0 \leq t \leq T$.

For each $0 \leq t \leq T$ let $H_t^\tau : M \rightarrow M$, $\tau \in \mathbb{R}$ be the solution flow to the vector field $X(\cdot)(h_t)$ so

$$\begin{cases} \frac{\partial}{\partial \tau} H_t^\tau(x) &= X(H_t^\tau(x))h_t \\ H_t^0(x) &= x. \end{cases} \quad (18)$$

Lemma 3.1 *The vector field U^h on $P\text{Diff}M$ has solution flow $\Phi_\tau : P\text{Diff}M \rightarrow P\text{Diff}M$, $\tau \in \mathbb{R}$ given by $\Phi_\tau(\phi)_t(x) = \phi_t(H_t^\tau(x))$.*

Proof. By left invariance we can suppose $\phi = e$. We then need only to observe that

$$\frac{\partial}{\partial \tau} H_t^\tau(x) = TH_t^\tau(X(x)h_t)$$

for each $0 \leq t \leq T$: a standard property of ordinary, time-independent dynamical systems which is seen by differentiating the identity

$$H_t^{\tau+\sigma} = H_t^\tau \circ H_t^\sigma(x)$$

with respect to σ at $\sigma = 0$. ■

C. In the case where h is random, with $h : \Omega \rightarrow L_0^{2,1}([0, T]; \mathbb{R}^d)$ adapted, we can use the same notation to obtain a variation of our stochastic flow $\{\xi_t : 0 \leq t \leq T\}$ on M generated by the vector field V^h , and given explicitly by

$$\xi^\tau = \Phi_\tau(\xi),$$

i.e.

$$\xi_t^\tau(x) = \xi_t(H_t^\tau(x)). \quad (19)$$

In particular

$$\frac{\partial}{\partial \tau} \xi_t^\tau(x) |_{\tau=0} = T\xi_t(X(x)h_t). \quad (20)$$

Using the structure of $C_x(M)$ as a C^∞ Banach manifold let $BC^1(C_x(M))$ be the space of C^1 maps $F : C_x(M) \rightarrow \mathbb{R}$ such that there is a constant $|dF|_\infty$ with

$$|dF(v)| \leq |dF|_\infty \sup_{0 \leq t \leq T} |v_t| \quad (21)$$

for all tangent vectors $v : [0, T] \rightarrow TM$ to $C_x(M)$. Set $V_t^{X(h)}(x) = T\xi_t(X(x)(h_t))$, which gives rise to a vector field along $\{\xi_t(x)\}$ on $C_x(M)$.

Proposition 3.2 Suppose $h : [0, T] \times \Omega \rightarrow T_x M$ is adapted, belongs to $L_0^{2,1}$ a.s. and such that $\mathbb{E} \left(\int_0^T |\dot{h}_s|^2 ds \right)^{\frac{1+\epsilon}{2}} < \infty$ for some $\epsilon > 0$. Then for each $x \in M$ the processes $\xi^\tau(x)$, $\tau \in \mathbb{R}$ have mutually equivalently laws \mathbb{P}_τ^x , $\tau \in \mathbb{R}$ on $C_x(M)$ with

$$\frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_\tau^x} = \exp \left\{ \int_0^T \left\langle X(\xi_s^\tau(x))^* T\xi_s \left(\frac{\partial}{\partial s} H_s^\tau(x) \right), dB_s \right\rangle - \frac{1}{2} \int_0^T |T\xi_s \left(\frac{\partial}{\partial s} H_s^\tau(x) \right)|^2 ds \right\}.$$

Moreover, for any $F \in BC^1(C_x(M))$,

$$\mathbb{E} dF(V^{X(h)}) = \mathbb{E} F(\xi) \int_0^T \left\langle X(\xi_s(x)) dB_s, V_s^{X(h)}(x) \right\rangle.$$

Proof. For the equivalent part note that $\{\xi_t^\tau : 0 \leq t \leq T\}$ satisfies the equation:

$$d\xi_t^\tau(x) = X(\xi_t^\tau(x)) \circ dB_t + A(\xi_t^\tau(x)) dt + T\xi_t \left(\frac{\partial}{\partial t} H_t^\tau(x) \right) dt.$$

A straightforward argument shows that

$$\int_0^T \left| X(\xi_s^\tau(x))^* T\xi_s \left(\frac{\partial}{\partial s} H_s^\tau(x) \right) \right|^2 < \infty, \quad a.s.$$

Therefore if we set

$$M_t^\tau = \int_0^t \left\langle X(\xi_s^\tau(x))^* T\xi_s \left(\frac{\partial}{\partial s} H_s^\tau(x) \right), dB_s \right\rangle,$$

then by the Girsanov-Maruyama theorem, P_τ^x is equivalent to P_0^x and

$$\frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_\tau^x} = e^{M_T^\tau - \frac{1}{2} \langle M \rangle_T^\tau}. \quad (22)$$

Consequently,

$$\mathbb{E} F(\xi^\tau(x)) = \mathbb{E} F(\xi(x)) \frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_\tau^x}.$$

Now suppose h . and $\int_0^T |\dot{h}_s|^2 ds$ are bounded on $[0, T] \times \Omega$. Differentiating with respect to τ at $\tau = 0$ and using (18) gives

$$\mathbb{E} dF(T\xi.(X(x)h.)) = \mathbb{E} F(\xi.(x)) \frac{\partial}{\partial \tau} \left(\frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_\tau^x} \right)_{\tau=0},$$

since $|dF|$ is bounded and $\sup_{0 \leq s \leq T} |T\xi_s| \in \cap_{1 \leq p \leq \infty} L^p$.

The second statement follows from differentiation of (22), using the fact that $\left(\frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_\tau^y}\right)_{\tau=0} = 1$ and $\frac{\partial}{\partial t} H_t^\tau(x)|_{\tau=0} = 0$:

$$\begin{aligned} \frac{\partial}{\partial \tau} \left(\frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_\tau^y} \right)_{\tau=0} &= \left(\frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_\tau^y} \right)_{\tau=0} \cdot \left[\left(\frac{\partial}{\partial \tau} M_T^\tau \right)_{\tau=0} - \frac{1}{2} \left(\frac{\partial}{\partial \tau} \langle M_T^\tau \rangle^2 \right)_{\tau=0} \right] \\ &= \int_0^T \left\langle X(\xi_s^\tau(x) dB_s, \frac{D}{\partial \tau} \left[T\xi_s \left(\frac{\partial}{\partial s} H_s^\tau(x) \right) \right] \right\rangle_{\tau=0} \\ &= \int_0^T \left\langle X(\xi_s(x) dB_s, T\xi_s \left(\frac{D}{\partial s} X(H_s^\tau(x)) h_s \right) \Big|_{\tau=0} \right\rangle \\ &= \int_0^T \left\langle X(\xi_s(x) dB_s, T\xi_s(X(x) \dot{h}_s)) \right\rangle. \end{aligned}$$

For general h take a sequence of bounded h_n which converges to h in $L^{\frac{1+\epsilon}{2}}(\Omega, L_0^{2,1}([0, T]))$ to finish the proof. See the proof of theorem 4.1. \blacksquare

The following is an analogue of Corollary 2.4: here $\tilde{\nabla}$ is any metric connection and \tilde{W}^Z is as in Corollary 2.4,

Theorem 3.3 *Let $F \in BC^1(C_x(M))$ and $h(\omega) \in L_0^{2,1}([0, T]; \mathbb{R}^n)$ a.s.. Suppose h is adapted to the filtration of $\{\mathcal{F}^x\}$ and such that $\mathbb{E} \left(\int_0^T |\dot{h}_s|^2 ds \right)^{\frac{1+\epsilon}{2}} < \infty$ for some $\epsilon > 0$. Then*

$$\mathbb{E} dF(\tilde{W}^Z(h.)) = \mathbb{E} F(\xi.(x)) \int_0^T \langle \tilde{W}_s^Z(\dot{h}_s), \tilde{\nabla}_s d\tilde{B}_s \rangle. \quad (23)$$

If $\tilde{\nabla}'$ is metric for some Riemannian metric, we can take $\epsilon = 0$.

4 Integration by parts for the free path space

It is easy to modify the proof of Proposition 3.2 to the case where $h(0) \neq 0$ and so obtain an integration by parts formula for the free path space $PM = \cup_{x \in M} P_x M$ with uniform topology and measure given by the Riemannian measure of M together with the laws of $\{\xi.(x) : x \in M\}$. In fact it is straightforward to generalize to the case of an x -dependent h . For this let $C^1(TM)$ be the space of C^1 vector fields on M with its usual topology:

Theorem 4.1 *Let $h : [0, T] \times \Omega \rightarrow C^1(TM)$ be a cadlag adapted process such that the $T_x M$ valued process $h.(x)$ has sample paths in $L^{2,1}([0, T]; T_x M)$*

for each $x \in M$ with $|h_0(\cdot)| + \sqrt{\int_0^t |\dot{h}_s(\cdot)|^2 ds}$ in $L^{1+\epsilon}(\Omega \times M; \mathbb{R})$ for some $\epsilon > 0$. Let F be in $BC^1(PM; \mathbb{R})$. Then

$$\begin{aligned} & \mathbb{E} \int_M dF(T_x \xi.(h.(\omega)(x))) dx \\ = & \mathbb{E} \int_M F(\xi.(x)) \left\{ -\operatorname{div} h_0(x) + \int_0^T \left\langle T \xi_s(\dot{h}_s(x)), X(\xi_s(x)) dB_s \right\rangle \right\} dx. \end{aligned} \quad (24)$$

Proof. Proceed as for Proposition 3.2 but with $X(x)h_t$ replaced by $h_t(x)$. In particular the definition (6) of H_t^τ becomes

$$\begin{aligned} \frac{\partial}{\partial \tau} H_t^\tau(x) &= h_t(H_t^\tau(x)) \\ H_t^0(x) &= x. \end{aligned}$$

while ξ_t^τ is defined by (19). However now $\xi_0^\tau(x) = \xi_0(H_0^\tau(x))$: the starting point is transported by the flow of $h_0(x)$.

We first assume $h.$ and $\int_0^t |\dot{h}_s|^2 ds$ are bounded on $\Omega \times M$. Then the Girsanov-Maruyama theorem gives us equivalence between the measures P_τ^x and $P_0^{H_0^\tau(x)}$ with

$$\int_M \mathbb{E} F(\xi^\tau(x)) dx = \int_M \mathbb{E} F(\xi.(H_0^\tau(x))) \frac{d\mathbb{P}_\tau^x}{d\mathbb{P}_0^{H_0^\tau(x)}} dx.$$

On differentiating this there is the extra term

$$\begin{aligned} & \int_M dF \left(T \xi. \left(\frac{\partial}{\partial \tau} H_0^\tau(x) \Big|_{\tau=0} \right) \right) \\ = & \int_M dF(T_x \xi.(h_0(x))) dx \\ = & \int_M d_x(F \circ \xi.)(h_0(x)) dx \end{aligned}$$

where $d_x(F \circ \xi.)$ refers to the derivative in M of $F \circ \xi. : M \times \Omega \rightarrow \mathbb{R}$. Now apply the classical Stokes theorem on M to get:

$$\begin{aligned} & \mathbb{E} \int_M dF(T_x \xi.(h.(\omega)(x))) dx \\ = & \mathbb{E} \int_M F(\xi.(x)) \left\{ -\operatorname{div} h_0(x) + \int_0^T \left\langle T_x \xi_s(\dot{h}_s(x)), X(\xi_s(x)) dB_s \right\rangle \right\} dx. \end{aligned}$$

For general h let τ_R be the first exit time of $\|h\|_{C^1} + \int_0^\cdot |h_s(x)|^2 ds$ from $[0, R)$. Set $h_t^R(x) = h_{t \wedge \tau_R}(x) \chi_{\{\|h_0\|_{C^1} < R\}}$. We have:

$$\begin{aligned} & \mathbb{E} \int_M dF(T_x \xi.(h^R(\omega)(x))) dx \\ = & \mathbb{E} \chi_{\{\|h_0\|_{C^1} < R\}} \int_M F(\xi.(x)) \left\{ -\operatorname{div} h_0(x) + \int_0^{T \wedge \tau_R} \langle T_x \xi_s(\dot{h}_s(x)), X(\xi_s(x)) dB_s \rangle \right\} dx. \end{aligned}$$

Now let $R \rightarrow \infty$. The left hand side converges to $\mathbb{E} \int_M dF(T\xi.(h(\omega)(x))) dx$ since

$$|dF(T\xi.(h^R(\omega)(-)))| \leq \tilde{c} \sup_t |T\xi_t(\omega)| \sup_t |h_t(-, \omega)|$$

and $\sup_x \mathbb{E} (\sup_t |T\xi_t| \int_M \sup_t |h_t(x, \omega)| dx) < \infty$ from

$$\begin{aligned} \sup_t |h_t(x)| & \leq |h_0(\omega)| + \int_0^T |\dot{h}_s(\omega)| ds \\ & \leq |h_0(\omega)| + \sqrt{T} \sqrt{\int_0^T |\dot{h}_s(\omega)|^2 ds} \in L^{1+\epsilon}(\Omega \times M) \end{aligned}$$

Using Burkholder-Davies-Gundy inequality to justify the integration on the right hand side we see that it converges to the right hand side of (24). ■

Just as before the intrinsic formulae can be deduced using [ELJL95]:

Theorem 4.2 *Let F be in $BC^1(PM; \mathbb{R})$ and h be as in Theorem 4.1 but with $h.(x)$ adapted to the filtration of $\{\mathcal{F}^x\}$, and $\operatorname{div} h_0 \in L^1(\Omega \times M, \mathbb{R})$. Then for any metric connection $\tilde{\nabla}$ on M ,*

$$\begin{aligned} & \mathbb{E} \int_M dF(\tilde{W}^Z(h(\omega)(x))) dx \\ = & \mathbb{E} \int_M F(\xi.(x)) \left\{ -\operatorname{div} h_0(x) + \int_0^T \left\langle \tilde{W}_s^Z(\dot{h}_s(x)), \tilde{\nabla}_s d\tilde{B}_s \right\rangle \right\} dx. \end{aligned} \tag{25}$$

If furthermore $\tilde{\nabla}'$ is metric with respect to a Riemannian metric, we can take $\epsilon = 0$.

Proof. The proof is just as that of Theorem 3.3. ■

References

- [Aid] S. Aida. On the irreducibility of certain Dirichlet forms on loop spaces over compact homogeneous spaces. To appear in 'New Trends in stochastic Analysis', Proc. Taniguchi Symposium, Sept. 1995, Charingworth, ed. K. D. Elworthy and S. Kusuoka, I. Shigekawa, World Scientific Press.
- [AM] H. Airault and P. Malliavin. Integration by parts formulas and dilation vector fields on elliptic probability spaces. Institut Mittag-Leffler preprints No. 24, 1994/95.
- [Bis81] J. M. Bismut. Martingales, the Malliavin calculus and harmonic theorems. In D. Williams, editor, *Stochastic Integrals, Lecture Notes in Maths. 851*, pages 85–109. Springer-Verlag, 1981.
- [Bis84] J. M. Bismut. *Large deviations and the Malliavin calculus. Progress in Math. 45*. Birkhäuser, 1984.
- [CM] A.-B. Cruzeiro and P. Malliavin. Curvatures of path spaces and stochastic analysis. Institut Mittag-Leffler preprints No. 16, 1994/95.
- [Dri92] B. Driver. A Cameron-Martin type quasi-invariance theorem for Brownian motion on a compact Riemannian manifold. *J. Funct. Anal.*, 100:272–377, 1992.
- [Dri95a] B. Driver. The Lie bracket of adapted vector fields on Wiener spaces. Preprint, 1995.
- [Dri95b] Bruce K. Driver. Towards calculus and geometry on path spaces. In *Stochastic Analysis: AMS Proceedings of symposium in pure Math. Series*, pages 423–426. AMS. Providence, Rhode Island, 1995.
- [EL94] K.D. Elworthy and Xue-Mei Li. Formulae for the derivatives of heat semigroups. *J. Funct. Anal.*, 125(1):252–286, 1994.
- [ELJL95] K. D. Elworthy, Yves Le Jan, and Xue-Mei Li. Concerning the geometry of stochastic differential equations and stochastic flows. To appear in 'New Trends in stochastic Analysis', Proc. Taniguchi Symposium, Sept. 1995, Charingworth, ed. K. D. Elworthy and S. Kusuoka, I. Shigekawa, World Scientific Press, 1995.
- [Elw82] K.D. Elworthy. *Stochastic Differential Equations on Manifolds*. Lecture Notes Series 70, Cambridge University Press, 1982.

- [Elw92] K. D. Elworthy. Stochastic flows on Riemannian manifolds. In M. A. Pinsky and V. Wihstutz, editors, *Diffusion processes and related problems in analysis, volume II. Birkhauser Progress in Probability*, pages 37–72. Birkhauser, Boston, 1992.
- [EM70] D. G. Ebin and J. Marsden. Groups of diffeomorphisms and the motion of an incompressible fluid. *Ann. of Math.*, 92(1):102–163, 1970.
- [ES95] O. Enchev and D.W. Stroock. Towards a Riemannian geometry on the path space over a Riemannian manifold. *J. Funct. Anal.*, 134(2):392–416, 1995.
- [EY93] K. D. Elworthy and M. Yor. Conditional expectations for derivatives of certain stochastic flows. In J. Azéma, P.A. Meyer, and M. Yor, editors, *Sem. de Prob. XXVII. Lecture Notes in Maths. 1557*, pages 159–172. Springer-Verlag, 1993.
- [FM93] S. Fang and P. Malliavin. Stochastic analysis on the path spaces of a Riemannian manifold. *J. Funct. Anal.*, 118:249–274, 1993.
- [Hsu95] E. Hsu. Inégalités de sobolev logarithmiques sur un espace de chemins. *C. R. Acad. Sci. Paris, t. 320. Série I.*, pages 1009–1012, 1995.
- [KN69] S. Kobayashi and K. Nomizu. *Foundations of differential geometry, Vol. II*. Interscience Publishers, 1969.
- [Li94] Xue-Mei Li. Stochastic differential equations on noncompact manifolds: moment stability and its topological consequences. *Probab. Theory Relat. Fields*, 100(4):417–428, 1994.
- [LN] R. Leandre and J. Norris. Integration by parts and Cameron-Martin formulas for the free-path space of a compact Riemannian manifold. Warwick Preprints: 6/1995.
- [Nel84] E. Nelson. *Quantum Fluctuations*. Princeton University Press, Princeton, 1984.
- [SZ] D. W. Stroock and O. Zeitouni. Variations on a theme by Bismut. Preprint.

Present address of Xue-Mei Li
 Mathematics Department, U-9, MSB 111, University of Connecticut,
 196 Auditorium Road, Storrs, Connecticut 06269, USA